

Erik Tijdsman⁽¹⁾, Kerstin Stahl⁽¹⁾, Sophie Bachmair⁽¹⁾ & Jamie Hannaford⁽²⁾


USA versus Europe

Introduction and Objectives

Climate zones or basin characteristics are often used to cluster and analyze drought characteristics.


- How representative are these groups regarding drought durations?
- What factors influence drought duration in the USA and Europe?
- How uniform are these factors for these two areas?
- Who has to deal with the longest droughts: **USA** or **Europe**?

Data and selection criteria



- Near natural streamflow data from: Hydro-Climatic Data Network dataset (**HCDN-2009**) and European Water Archive (**EWA**)
- Continuous and daily data availability between **1970-2010** and **1965-2005**
- Daily 20-percentile flow is never zero
- Precipitation and temperature from **PRISM** (800m resolution, via USGS) and **E-OBS** (0.25° resolution).
- Köppen-Geiger climates and aridity index are derived from these data


Drought duration identification



Testing influential factors





Approach 1: Single factors




- **Precipitation** decreases with increasing DDC classes. The differences are mostly significant, except for similarities between some of the neighboring classes, indicating a strong influence of this factor.
- **Elevation** shows a different pattern for the different areas. Where longer drought duration classes are located at higher elevations in the USA, it is the other way around for Europe. Longer drought duration classes have similar elevations and these elevations are significantly different from the other drought duration classes.

Similarity between areas? Aridity index USA vs. Europe



Approach 2: Climate classification factors



- **Köppen-Geiger** climates In the USA show an ordering of DDC, where non-seasonal warm summer climates have the shortest droughts, followed by non-seasonal hot summer climates and by the longest droughts for the seasonal and cold summer climates. However, these differences are often not significant, only between the non-seasonal warm summer climates and the rest. For Europe, only the Cfc climate shows significantly shorter droughts.
- **The aridity index** shows a clear ordering of DDC from the most arid to the least arid groups. These differences are mostly significant, except some similarities between neighboring classes.

Conclusions

- The USA have longer droughts and different factors influence (the difference in) drought duration.
- Absolute precipitation has a similar and significant influence on DDC for both areas.
- DDC grouped by Köppen-Geiger climate classification show the influence of seasonality and hot/cold summer climates on DDC. However, differences are mostly not significant.
- The aridity index is a good proxy for drought durations on both continents. However, they are not directly comparable between the two areas.